



FLORIDA SOLAR ENERGY CENTER

A Research Institute of the University of Central Florida

Hydrogen Production via Photolytic Oxidation of Aqueous Sodium Sulfite Solutions

*Cunping Huang, **Clovis A. Linkous**,
Olawale Adebisi & Ali T-Raissi*

Hydrogen 2008 – Materials Innovations in an Emerging
Hydrogen Economy

Cocoa Beach, Florida, USA
February 27, 2008



Background

- ❖ SO₂ is a criteria air pollutant that can cause respiratory & other problems as an acid gas
- ❖ SO₂ emissions occur both naturally (20%) & anthropogenically (80%)
- ❖ Natural sources include: geothermal (e.g. volcanic), oceanic, vegetative & land emissions.



Anthropogenic Sources of SO₂ Emissions

- ❖ Combustion of high-sulfur-containing fossil fuels
- ❖ Sulfuric acid & ammonium sulfate plants
- ❖ Power plants using coal, crude oil & crude oil-based fuel oil.




Major Global Anthropogenic Sources of SO₂ Emissions

Emission Source	Emission (%)
Electric utility	69.7
Industrial fuel combustion	13.6
Metal processing	3.8
Transportation	3.5
Others	9.4

Source: Schnelle, K.B., and Brown, C.A., Control of So_x, In Air Pollution Control Technology Handbook, ed. Kreith, F., CRC Press, Boca Raton, FL, 257, 2002



SO₂ is both a Pollutant & a Resource



Generation of SO₂ in 500 MW coal fired power plants can produce huge amounts of SO₂ that can be used for the production of H₂ as well as fertilizers.



Objectives

- ❖ Develop an innovative process for utilizing SO_2 in flue gas for the production of hydrogen
- ❖ Explore chemistry & chemical engineering aspects of SO_2 utilization
- ❖ Investigate effects of reaction conditions on the hydrogen production rate.



Flue Gas Treatment and H₂ Production

Conventional process:



FSEC Approach:




Photooxidation:






Experimental Setup for SO_2 Treatment






Exp. Setup for H_2 Production from Aqueous Na_2SO_3 Solution






Hydrogen Production





Kinetics of H_2 Production via Photo-oxidation of Na_2SO_3






Material Balance

Ionic Species	Initial (mmol)	Final (mmol)	Diff. (mmol)
SO_3^{2-}	63.41	0.00	63.41
SO_4^{2-}	2.40	63.59	61.19
Gas Produced	Theoretical (mL)	Exp. (mL)	Diff. (mL)
H_2	1550	1390	160




Effect of Solution pH on H_2 Production






Effect of Solution pH on H₂ Production (cont'd)






Concentration Effect on H_2 Production Rate (at pH = 9.70)





Concentration Effect on H_2 Production Rate (at pH = 7.55)





Summary

Solution	Conc. 0.025 M	pH 9.92	Conc. 0.05 M	pH 9.70	Conc. 0.025 M	pH 7.53	Conc. 0.05 M	pH 7.59
H ₂ prod rate	0.74 mL/min		1.40 mL/min		1.91 mL/min		1.91 mL/min	

At pH = 9.95, H₂ production rate increases with an increase in the concentration of the sulfite.

At pH = 7.55, H₂ production rate is independent of the sulfite concentration.



Conclusions

- ❖ A novel approach for utilizing SO_2 in flue gas for hydrogen production has been developed
- ❖ Photolytic H_2 production from aqueous Na_2SO_3 solutions is a clean and efficient process
- ❖ Experimental data indicate that SO_3^{2-} can be fully converted into SO_4^{2-}
- ❖ FSEC process requires no catalysts, reducing the process capital & operating costs.



Future Work

- ❖ Investigate effects of other flue gases (e.g. NO_x , CO_2) on the photolytic production of hydrogen
- ❖ Investigate effects of metal catalysts in enhancing the photolytic hydrogen production from SO_2
- ❖ Photoreactor design considerations
- ❖ Process design & optimization.



Acknowledgment

- ❖ National Aeronautics and Space Administration (NASA) - Glenn Research Center (GRC) under contract No. NAG3-2751.